



Melanie Muhlstein, M.D.

5310 Hampton Place, Suite 1 | Saginaw, MI 48604

989-321-2655 | Fax: 989-321-2661

Patient Care Instructions Post Intravitreal Injection

Please follow these instructions after eye injections:

- Do not rub your eye after treatment
- For relief of mild to moderate pain or irritation, an over-the-counter pain reliever (i.e. Tylenol) or cool compress/ice pack can be used
- A new bottle of artificial tears/lubricating drops can be used
- No swimming or contact lens wear for 2 days
- It is okay to shower and wash your face. Keep eyes closed to avoid dirty water in the eyes.
- Please continue medicated eye drops as previously prescribed

The following symptoms are **COMMON** after successful eye injections for a day or so:

- Blurry vision
- Tearing
- Mild floaters or bubbles in your vision
- Mild to moderate discomfort or irritation
- Gritty, "Eyelash in the eye" sensation
- Redness (sometimes this can cover the entire white part of the eye and last a few weeks)
- Bloody tears for 1 - 2 days after treatment

The following symptoms are **NOT NORMAL** after eye injections, and you should call our office immediately if you experience:

- Severe, constant pain
- Worsening pain after the first day
- Significant decrease in vision
- Severe increase in floaters
- Curtain or veil in your vision

Please contact our office with any questions or concerns. Follow-up care is very important. Be sure to keep your scheduled appointment.